

Dirichlet L -Functions and Character Power Sums

TOM M. APOSTOL

*Department of Mathematics, California Institute of Technology,
Pasadena, California 91109*

Communicated by S. Chowla

Received October 9, 1969; revised November 17, 1969

A new representation for Dirichlet L -functions $L(s, \chi)$, valid for primitive characters χ modulo k and all complex s , is given in terms of the function $F(x, s)$ defined for real x and $R(s) > 1$ by the series $\sum_{n=1}^{\infty} e^{2n\pi ix}/n^s$. Evaluation of $L(s, \chi)$ for negative integer s leads to a class of identities relating m th power moments $\sum_{r=1}^{k-1} \chi(r)r^m$ with finite cotangent power sums. Special emphasis is given to the quadratic character $\chi(n) = (n|p)$, p an odd prime. A new proof of the functional equation for L -functions is also given.

1. INTRODUCTION

Let k be a positive integer and let χ be any character modulo k . The L -function $L(s, \chi)$, defined for $R(s) > 1$ by the Dirichlet series

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s},$$

can be expressed in terms of the Hurwitz zeta function $\zeta(s, a)$ by means of the formula

$$L(s, \chi) = k^{-s} \sum_{h=1}^{k-1} \chi(h) \zeta\left(s, \frac{h}{k}\right), \quad (1)$$

which is valid for all complex s [6, p. 113]. Here $\zeta(s, a)$ is the analytic continuation of the series

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s}, \quad R(s) > 1, \quad 0 < a \leq 1.$$

In this paper we derive an analogous representation for $L(s, \chi)$, which we

believe to be new, in terms of the function $\varphi(z, s)$ which is defined by the power-Dirichlet series

$$\varphi(z, s) = \sum_{n=1}^{\infty} \frac{z^n}{n^s}. \quad (2)$$

This representation, stated below in Theorem 1, is valid only for primitive characters.

The series in (2) converges for all s if $|z| < 1$, for $R(s) > 0$ if $|z| = 1$, $z \neq 1$, and for $R(s) > 1$ if $z = 1$. The function $\varphi(z, s)$ has been studied by many writers, including Lambert, Legendre, Abel, Kummer, Appell, Lerch, Jonquière, Lindelöf, Wirtinger, Truesdell [7], and others. When $z = 1$, $\varphi(z, s)$ is the Riemann zeta function, $\zeta(s)$.

By use of the classic method of Riemann, $\varphi(z, s)$ can be extended to the whole s -plane by means of the contour integral [2, p. 28], [4, p. 144]

$$I(z, s) = \frac{1}{2\pi i} \int_C \frac{t^{s-1} z e^t}{1 - z e^t} dt,$$

where C is a loop in the t -plane which begins at $-\infty$, encircles the origin once in the positive direction, and returns to $-\infty$. If the z -plane is cut from 1 to ∞ along the positive real axis, then for every fixed z in the cut plane, $I(z, s)$ is an entire function of s , and we have

$$\varphi(z, s) = \Gamma(1-s) I(z, s).$$

We are interested here in values of z on the unit circle $|z| = 1$, with $z \neq 1$. We let $z = e^{2\pi i x}$, where x is real, and we define $F(x, s) = \varphi(e^{2\pi i x}, s)$. Clearly, $F(x, s)$ is periodic in x with period 1. For $x \neq \text{integer}$ and $R(s) > 0$ we have the series representation

$$F(x, s) = \sum_{n=1}^{\infty} \frac{e^{2\pi i n x}}{n^s}. \quad (3)$$

When x is an integer we have $F(x, s) = \zeta(s)$.

In Section 2 we express $L(s, \chi)$ in terms of $F(x, s)$, and in Section 3 we use this representation to evaluate $L(s, \chi)$ when s is a negative integer or 0. This leads to a class of identities relating m th power moments

$$M_m(\chi) = \sum_{r=1}^{k-1} \chi(r) r^m \quad (4)$$

with cotangent power sums

$$T_m(\chi) = \sum_{r=1}^{k-1} \chi(r) \cot^m(\pi r/k) \quad (5)$$

for $m = 1, 2, 3, \dots$.

In a study on quadratic residues, Lebesgue [5, p. 233] obtained the formula

$$\sum_{r=1}^{p-1} r(r|p) = -\frac{1}{2} \sqrt{p} \sum_{r=1}^{p-1} \cot(\pi r^2/p) \quad (6)$$

for primes $p \equiv 3 \pmod{4}$, where $(r|p)$ is the quadratic character symbol of Legendre. In Section 4 we show that Lebesgue's formula is a very special case of the formulas in Section 3, obtained by choosing $m = 1$ and taking $\chi(n)$ to be the quadratic character $(n|p)$.

In Section 5 we again take $\chi(n) = (n|p)$, where p is an odd prime, and we obtain an alternate representation of $L(s, \chi)$ in terms of both $F(x, s)$ and $\zeta(s)$. This is used in Section 6 to evaluate $L(s, \chi)$ for negative integer values of s , giving us another set of identities relating the m th power moments

$$M_m = \sum_{r=1}^{p-1} r^m (r|p)$$

with the cotangent power sums

$$S_m = \sum_{r=1}^{p-1} \cot^m(\pi r^2/p).$$

For $p \equiv 3 \pmod{4}$ these turn out to be the same identities obtained in Section 4, but for $p \equiv 1 \pmod{4}$ they are different from those of Section 4.

Finally, in Section 7 we use the representation of Theorem 1 to give a new and short proof of the functional equation for L -functions.

2. A REPRESENTATION THEOREM FOR L -FUNCTIONS

For any character χ modulo k let $G(m, \chi)$ denote the Gauss sum

$$G(m, \chi) = \sum_{h=1}^{k-1} \chi(h) e^{2\pi i m h/k},$$

and let $G(\chi) = G(1, \chi)$. Then we have:

THEOREM 1. *If χ is a primitive character modulo k , then for all complex s we have the representation*

$$G(\bar{\chi}) L(s, \chi) = \sum_{h=1}^{k-1} \bar{\chi}(h) F\left(\frac{h}{k}, s\right). \quad (7)$$

Proof. We prove the result first for $R(s) > 1$ and then use analytic continuation to extend it for all s .

Starting with Eq. (3), we take $x = h/k$, multiply both members by $\bar{\chi}(h)$, and sum on h to obtain

$$\sum_{h=1}^{k-1} \bar{\chi}(h) F\left(\frac{h}{k}, s\right) = \sum_{n=1}^{\infty} n^{-s} \sum_{h=1}^{k-1} \bar{\chi}(h) e^{2\pi i n h/k} = \sum_{n=1}^{\infty} n^{-s} G(n, \bar{\chi}).$$

But since χ is a primitive character we have the factorization property [1, p. 312]

$$G(n, \bar{\chi}) = \chi(n) G(1, \bar{\chi}) = \chi(n) G(\bar{\chi}),$$

which holds for every n . Hence we obtain

$$\sum_{n=1}^{\infty} n^{-s} G(n, \bar{\chi}) = G(\bar{\chi}) \sum_{n=1}^{\infty} \chi(n) n^{-s} = G(\bar{\chi}) L(s, \chi),$$

which proves (7).

3. COTANGENT IDENTITIES FOR CHARACTER POWER SUMS

The two representations for $L(s, \chi)$ in (1) and (7) give us the identity

$$G(\bar{\chi}) k^{-s} \sum_{r=1}^{k-1} \chi(r) \zeta\left(s, \frac{r}{k}\right) = \sum_{h=1}^{k-1} \bar{\chi}(h) F\left(\frac{h}{k}, s\right), \quad (8)$$

valid for all s . In this formula we substitute special values of s for which the functions $\zeta(s, a)$ and $F(x, s)$ can be evaluated explicitly. For example, if $s = 0$ we have [2, p. 27]

$$\zeta(0, a) = \frac{1}{2} - a.$$

From (2) we find $\varphi(z, 0) = z/(1 - z)$ if $|z| \leq 1, z \neq 1$. Taking $z = e^{2\pi i x}$, we obtain

$$F(x, 0) = -\frac{1}{2} + \frac{i}{2} \cot \pi x \quad (9)$$

if x is not an integer. Putting these values in (8) and using the relation $\sum_{r=1}^{k-1} \chi(r) = 0$ we obtain the identity

$$G(\bar{\chi}) \sum_{r=1}^{k-1} r \chi(r) = -\frac{1}{2} i k \sum_{h=1}^{k-1} \bar{\chi}(h) \cot(\pi h/k).$$

In the notation of (4) and (5) we have proved that

$$G(\bar{\chi}) M_1(\chi) = -\frac{1}{2} ik T_1(\bar{\chi}). \quad (10)$$

The functions $\zeta(s, a)$ and $F(x, s)$ can also be evaluated when s is a negative integer. If $n \geq 1$ we have [2, p. 27]

$$\zeta(-n, a) = -B_{n+1}(a)/(n+1), \quad (11)$$

where $B_{n+1}(a)$ is a Bernoulli polynomial. The function values $F(x, -n)$ can be calculated recursively from the differential-difference equation

$$F(x, s-1) = \frac{1}{2\pi i} \frac{\partial F(x, s)}{\partial x}. \quad (12)$$

This follows from (3) for $R(s) > 2$, and by analytic continuation for all s . Using (12) repeatedly with the initial value $F(x, 0)$ in (9) we find

$$F(x, -n) = \frac{1}{(2\pi i)^n} \frac{d^n}{dx^n} \left(\frac{i}{2} \cot \pi x \right). \quad (13)$$

The first few values are

$$F(x, -1) = \frac{-1}{4} (1 + \cot^2 \pi x),$$

$$F(x, -2) = \frac{-i}{8} (2 \cot \pi x + 2 \cot^3 \pi x),$$

$$F(x, -3) = \frac{1}{16} (2 + 8 \cot^2 \pi x + 6 \cot^4 \pi x),$$

$$F(x, -4) = \frac{i}{32} (16 \cot \pi x + 40 \cot^3 \pi x + 24 \cot^5 \pi x).$$

It is easy to prove by induction that for any $n \geq 1$, $F(x, -n)$ is a polynomial in $\cot \pi x$ of degree $n+1$ given by

$$F(x, -n) = \frac{i^{n+1}}{2^{n+1}} \sum_{k=0}^{n+1} a_{n,k} \cot^k \pi x.$$

The coefficients $a_{n,k}$ are nonnegative integers which can be readily calculated from the following recursion formulas:

$$a_{0,0} = 0, \quad a_{0,1} = 1,$$

$$a_{n,0} = a_{n-1,1} \quad \text{for } n \geq 1,$$

$$a_{n,k} = (k-1) a_{n-1,k-1} + (k+1) a_{n-1,k+1} \quad \text{for } 1 \leq k \leq n,$$

$$a_{n,n+1} = n! \quad \text{for } n \geq 1.$$

The coefficients for $n \leq 6$ are listed in Table I.

TABLE I

n	$a_{n,0}$	$a_{n,1}$	$a_{n,2}$	$a_{n,3}$	$a_{n,4}$	$a_{n,5}$	$a_{n,6}$	$a_{n,7}$
1	1	0	1					
2	0	2	0	2				
3	2	0	8	0	6			
4	0	16	0	40	0	24		
5	16	0	136	0	240	0	120	
6	0	272	0	1232	0	1680	0	720

Putting $s = -n$ in Theorem 1 and multiplying by $-(n+1)$ we find

$$k^n G(\bar{\chi}) \sum_{r=1}^{k-1} \chi(r) B_{n+1}\left(\frac{r}{k}\right) = -(n+1) \sum_{h=1}^{k-1} \bar{\chi}(h) F\left(\frac{h}{k}, -n\right), \quad (14)$$

for $n = 1, 2, 3, \dots$. Taking $n = 1$ and using the formula

$$B_2(x) = x^2 - x + \frac{1}{6}$$

we are led to the relation

$$G(\bar{\chi}) M_2(\chi) = kG(\bar{\chi}) M_1(\chi) + \frac{1}{2}kT_2(\bar{\chi}).$$

Using (10) to express $G(\bar{\chi}) M_1(\chi)$ in terms of $T_1(\bar{\chi})$ we find

$$G(\bar{\chi}) M_2(\chi) = -\frac{1}{2}ik^2T_1(\bar{\chi}) + \frac{1}{2}kT_2(\bar{\chi}). \quad (15)$$

When $n = 2$ we have $B_3(x) = x^3 - \frac{3}{2}x^2 + \frac{1}{2}x$. This leads to the formula

$$G(\bar{\chi}) M_3(\chi) = \frac{3}{2}kG(\bar{\chi}) M_2(\chi) - \frac{1}{2}k^2G(\bar{\chi}) M_1(\chi) + \frac{3}{4}ik\{T_1(\bar{\chi}) + T_3(\bar{\chi})\}.$$

Using (10) and (15) we find

$$G(\bar{\chi}) M_3(\chi) = i(\frac{3}{4}k - \frac{1}{2}k^3) T_1(\bar{\chi}) + \frac{3}{4}k^2T_2(\bar{\chi}) + \frac{3}{4}ikT_3(\bar{\chi}). \quad (16)$$

When $n = 3$ we have $B_4(x) = x^4 - 2x^3 + x^2 - 1/30$. When this is used in (14) we obtain

$$G(\bar{\chi}) M_4(\chi) = 2kG(\bar{\chi}) M_3(\chi) - k^2G(\bar{\chi}) M_2(\chi) - 2kT_2(\bar{\chi}) - \frac{3}{2}kT_4(\bar{\chi}).$$

Using (15) and (16) we find

$$\begin{aligned} G(\bar{\chi}) M_4(\chi) \\ = \frac{1}{2}ik^2(3 - k^2) T_1(\bar{\chi}) + k(k^2 - 2) T_2(\bar{\chi}) + \frac{3}{2}ik^2T_3(\bar{\chi}) - \frac{3}{2}kT_4(\bar{\chi}). \end{aligned} \quad (17)$$

It is clear that, in general, the number $G(\bar{\chi}) M_m(\chi)$ can be expressed as a linear combination of the cotangent power sums $T_1(\bar{\chi}), \dots, T_m(\bar{\chi})$, the coefficients being polynomials in k .

4. QUADRATIC RESIDUE POWER SUMS

This section considers some special cases of the formulas in Section 3. We take $k = p$, where p is an odd prime, and let $\chi(n) = (n | p)$, the quadratic character modulo p , where it is understood that $(n | p) = 0$ if $n \equiv 0 \pmod{p}$. For this choice of χ the Gauss sum $G(\chi)$ is given by [1, p. 315]

$$G(\chi) = \sum_{r=1}^{p-1} (r | p) e^{2\pi i r/p} = \begin{cases} \sqrt{p} & \text{if } p \equiv 1 \pmod{4} \\ i\sqrt{p} & \text{if } p \equiv 3 \pmod{4}. \end{cases} \quad (18)$$

Therefore the formulas in Section 3 split naturally into two cases, $p \equiv 1$ and $p \equiv 3 \pmod{4}$. We write M_m for $M_m(\chi)$ and T_m for $T_m(\chi)$ so that

$$M_m = \sum_{r=1}^{p-1} (r | p) r^m, \quad T_m = \sum_{h=1}^{p-1} (h | p) \cot^m(\pi h/p).$$

Both sums M_m and T_m are real-valued. Using (18), and equating real and imaginary parts in Eq. (10), we obtain the pair of formulas

$$M_1 = T_1 = 0 \quad \text{if} \quad p \equiv 1 \pmod{4} \quad (19)$$

and

$$M_1 = -\frac{1}{2} \sqrt{p} T_1 \quad \text{if } p \equiv 3 \pmod{4}. \quad (20)$$

It is easy to show that (20) contains Lebesgue's formula (6). Since the cotangent is an odd function with period π we have

$$\sum_{r=1}^{p-1} \cot(\pi r/p) = 0.$$

Therefore we have

$$\begin{aligned} T_1 &= \sum_{h=1}^{p-1} (h | p) \cot(\pi h/p) = \sum_{h=1}^{p-1} \{(h | p) + 1\} \cot(\pi h/p) \\ &= 2 \sum_{\substack{h=1 \\ (h|p)=1}}^{p-1} \cot(\pi h/p) = \sum_{r=1}^{p-1} \cot(\pi r^2/p), \end{aligned} \quad (21)$$

so Eq. (20) reduces to Lebesgue's formula (6).

The identities in Section 3 corresponding to $m = 2, 3, 4$ lead to the following pairs of formulas:

$$M_2 = \begin{cases} \frac{1}{2} \sqrt{p} T_2 & \text{if } p \equiv 1 \pmod{4} \\ -\frac{1}{2} p \sqrt{p} T_1 & \text{if } p \equiv 3 \pmod{4}, \end{cases} \quad (22)$$

$$M_3 = \begin{cases} \frac{3}{4} p \sqrt{p} T_2 & \text{if } p \equiv 1 \pmod{4} \\ -\frac{1}{4} \sqrt{p} (2p^2 - 3) T_1 + \frac{3}{4} \sqrt{p} T_3 & \text{if } p \equiv 3 \pmod{4}, \end{cases} \quad (23)$$

$$M_4 = \begin{cases} \sqrt{p} (p^2 - 2) T_2 - \frac{3}{2} \sqrt{p} T_4 & \text{if } p \equiv 1 \pmod{4} \\ -\frac{1}{2} \sqrt{p} p (p^2 - 3) T_1 + \frac{3}{2} p \sqrt{p} T_3 & \text{if } p \equiv 3 \pmod{4}. \end{cases} \quad (24)$$

As a by-product we also obtain the relation $T_{2m+1} = 0$ for $p \equiv 1 \pmod{4}$, and $T_{2m} = 0$ for $p \equiv 3 \pmod{4}$.

5. ALTERNATE REPRESENTATION FOR $L(s, \chi)$ WHEN $\chi(n) = (n | p)$.

Let p be an odd prime, let $\chi(n) = (n | p)$, let $G = G(1, \chi)$, and let $L(s) = L(s, \chi)$. Then we have:

THEOREM 2. *For all complex s we have the representation*

$$GL(s) = \sum_{h=1}^{p-1} F\left(\frac{h^2}{p}, s\right) + (1 - p^{1-s}) \zeta(s). \quad (25)$$

Proof. Taking $k = p$ and $\chi(n) = (n | p)$ in Theorem 1 we have

$$GL(s) = \sum_{r=1}^{p-1} (r | p) F\left(\frac{r}{p}, s\right) = \sum_{\substack{r=1 \\ (r|p)=1}}^{p-1} F\left(\frac{r}{p}, s\right) - \sum_{\substack{r=1 \\ (r|p)=-1}}^{p-1} F\left(\frac{r}{p}, s\right). \quad (26)$$

Using (3) we have, for $R(s) > 1$,

$$\sum_{r=1}^p F\left(\frac{r}{p}, s\right) = \sum_{n=1}^{\infty} n^{-s} \sum_{r=1}^p e^{2\pi i n r / p} = p \sum_{\substack{n=1 \\ n \equiv 0 \pmod{p}}}^{\infty} n^{-s} = p^{1-s} \zeta(s).$$

Therefore, by analytic continuation, for all s we have

$$p^{1-s} \zeta(s) = \sum_{r=1}^p F\left(\frac{r}{p}, s\right) = \sum_{\substack{r=1 \\ (r|p)=1}}^{p-1} F\left(\frac{r}{p}, s\right) + \sum_{\substack{r=1 \\ (r|p)=-1}}^{p-1} F\left(\frac{r}{p}, s\right) + F(1, s). \quad (27)$$

Adding (26) and (27) we obtain

$$\begin{aligned} GL(s) &= 2 \sum_{\substack{r=1 \\ (r|p)=1}}^{p-1} F\left(\frac{r}{p}, s\right) + (1 - p^{1-s}) \zeta(s) \\ &= \sum_{h=1}^{p-1} F\left(\frac{h^2}{p}, s\right) + (1 - p^{1-s}) \zeta(s), \end{aligned}$$

which proves (25).

6. FURTHER COTANGENT IDENTITIES FOR QUADRATIC CHARACTERS

Using Eq. (1) we can rewrite the identity of Theorem 2 in the form

$$Gp^{-s} \sum_{r=1}^{p-1} (r|p) \zeta\left(s, \frac{r}{p}\right) = \sum_{h=1}^{p-1} F\left(\frac{h^2}{p}, s\right) + (1 - p^{1-s}) \zeta(s).$$

In this formula we substitute $s = -n$ ($n = 0, 1, 2, 3, \dots$), using (11), and the relation [2, p. 34]

$$\zeta(-n) = -B_{n+1}/(n+1),$$

where B_{n+1} is a Bernoulli number. This gives us

$$Gp^n \sum_{r=1}^{p-1} (r|p) B_{n+1} \left(\frac{r}{p}\right) = (p^{n+1} - 1) B_{n+1} - (n+1) \sum_{h=1}^{p-1} F\left(\frac{h^2}{p}, -n\right). \quad (28)$$

Now we take $n = 0, 1, 2, 3$ and proceed as in Section 4 (omitting details) to obtain the following formulas:

$$GM_1 = -\frac{1}{2}ipS_1, \quad (29)$$

$$GM_2 = \frac{1}{2}pS_2 - \frac{1}{6}p(p-1)(p-2) - \frac{1}{2}ip^2S_1, \quad (30)$$

$$GM_3 = -\frac{1}{4}p^2(p-1)(p-2) + i\left(\frac{3}{4}p - \frac{1}{2}p^3\right)S_1 + \frac{3}{4}p^2S_2 + \frac{3}{4}ipS_3, \quad (31)$$

$$\begin{aligned} GM_4 &= \frac{1}{30}p(p-1)(p-2)(7+3p-9p^2) + \frac{1}{2}ip^2(3-p^2)S_1 \\ &\quad + p(p^2-2)S_2 + \frac{3}{2}ip^2S_3 - \frac{3}{2}pS_4. \end{aligned} \quad (32)$$

Using the value of G in (18) we obtain the following pairs of formulas:

$$M_2 = \begin{cases} \frac{1}{2} \sqrt{p} S_2 - \frac{1}{6} \sqrt{p} (p-1)(p-2) & \text{if } p \equiv 1 \pmod{4} \\ -\frac{1}{2} p \sqrt{p} S_1 & \text{if } p \equiv 3 \pmod{4}, \end{cases} \quad (33)$$

$$M_3 = \begin{cases} \frac{3}{4} \sqrt{p} p S_2 - \frac{1}{4} \sqrt{p} p (p-1)(p-2) & \text{if } p \equiv 1 \pmod{4} \\ \frac{3}{4} \sqrt{p} S_3 - \frac{1}{4} \sqrt{p} (2p^2-3) S_1 & \text{if } p \equiv 3 \pmod{4}, \end{cases} \quad (34)$$

$$M_4 = \begin{cases} \frac{1}{36} \sqrt{p} (p-1)(p-2)(7+3p-9p^2) + \sqrt{p} (p^2-2) S_2 - \frac{3}{2} \sqrt{p} S_4 & \text{if } p \equiv 1 \pmod{4} \\ \frac{3}{2} \sqrt{p} p S_3 - \frac{1}{2} \sqrt{p} p (p^2-3) S_1 & \text{if } p \equiv 3 \pmod{4}. \end{cases}$$

We also obtain the relations

$$S_2 = \frac{1}{3}(p-1)(p-2) \quad \text{if } p \equiv 3 \pmod{4}$$

and

$$S_4 = \frac{1}{45}(p-1)(p-2)(p^2+3p-13) \quad \text{if } p \equiv 3 \pmod{4}.$$

By the same argument used in Eq. (21) to prove that $T_1 = S_1$ we see that the cotangent sums T_m and S_m are equal when m is odd. Therefore, for $p \equiv 3 \pmod{4}$ the above formulas for M_1, \dots, M_4 are the same as those obtained in Section 4. By comparing the formulas (33) through (35) with the corresponding results in Section 4 we also obtain the following cotangent identities, valid for primes $p \equiv 1 \pmod{4}$:

$$T_2 = S_2 - \frac{1}{3}(p-1)(p-2) \quad \text{if } p \equiv 1 \pmod{4}$$

and

$$T_4 = S_4 - \frac{1}{45}(p-1)(p-2)(p^2+3p-13) \quad \text{if } p \equiv 1 \pmod{4}.$$

7. THE FUNCTIONAL EQUATION FOR L -FUNCTIONS

In this section we give a simple proof of the functional equation for L -functions [6, p. 307].

THEOREM 3. *If χ is any primitive character modulo k , then for all complex s we have*

$$L(1-s, \chi) = (2\pi)^{-s} \Gamma(s) k^{s-1} \{e^{-is\pi/2} + \chi(-1) e^{is\pi/2}\} G(\chi) L(s, \bar{\chi}). \quad (36)$$

Proof. Our proof is based on the formula [2, p.31]

$$\zeta(1-s, x) = (2\pi)^{-s} \Gamma(s) \{e^{-is\pi/2} F(x, s) + e^{is\pi/2} F(-x, s)\}, \quad (37)$$

which is valid for $0 < x \leq 1$ and all complex s . This relation is usually attributed to Jonquière [4] who proved it in 1889. However, it is really an alternate form of a well-known formula of Hurwitz [3], proved in 1882, which states that [2, p. 26]

$$\zeta(s, x) = 2(2\pi)^{s-1} \Gamma(1-s) \sum_{n=1}^{\infty} \frac{\sin(2n\pi x + \pi s/2)}{n^{1-s}} \quad (38)$$

if $R(s) < 0$ and $0 < x \leq 1$. If we replace s by $1-s$ and express $\sin(2n\pi x + \pi s/2)$ in terms of exponentials, Eq. (38) takes the form

$$\zeta(1-s, x) = (2\pi)^{-s} \Gamma(s) \left\{ e^{-is\pi/2} \sum_{n=1}^{\infty} \frac{e^{2\pi i n x}}{n^s} + e^{is\pi/2} \sum_{n=1}^{\infty} \frac{e^{-2\pi i n x}}{n^s} \right\}$$

provided $R(s) > 1$. This clearly implies (37) for $R(s) > 1$, and the result holds for all s by analytic continuation.

To derive the functional equation (36) from (37) we simply take $x = h/k$ in (37), then multiply both members by $\chi(h)$ and sum on h . This gives us

$$\begin{aligned} & \sum_{h=1}^{k-1} \chi(h) \zeta\left(1-s, \frac{h}{k}\right) \\ &= \frac{\Gamma(s)}{(2\pi)^s} \left\{ e^{-is\pi/2} \sum_{h=1}^{k-1} \chi(h) F\left(\frac{h}{k}, s\right) + e^{is\pi/2} \sum_{h=1}^{k-1} \chi(h) F\left(\frac{-h}{k}, s\right) \right\} \\ &= \frac{\Gamma(s)}{(2\pi)^s} \{e^{-is\pi/2} + \chi(-1) e^{is\pi/2}\} \sum_{h=1}^{k-1} \chi(h) F\left(\frac{h}{k}, s\right), \end{aligned} \quad (39)$$

since

$$\begin{aligned} \sum_{h \bmod k} \chi(h) F\left(\frac{-h}{k}, s\right) &= \chi(-1) \sum_{h \bmod k} \chi(-h) F\left(\frac{-h}{k}, s\right) \\ &= \chi(-1) \sum_{h \bmod k} \chi(h) F\left(\frac{h}{k}, s\right). \end{aligned}$$

Now we multiply both members of (39) by k^{s-1} , then use (1) on the left and Theorem 1 on the right to obtain (36).

REFERENCES

1. R. AYOUB, "An Introduction to the Analytic Theory of Numbers," Mathematical Surveys, Number 10, American Mathematical Society, Providence, R. I., 1963.
2. A. ERDÉLYI (editor), "Higher Transcendental Functions," vol. I, Bateman Manuscript Project, McGraw-Hill, New York, 1953.
3. A. HURWITZ, Einige Eigenschaften der Dirichlet'schen Funktionen $F(s) = \sum (D|n)n^{-s}$, die bei der Bestimmung der Klassenanzahlen binärer quadratischer Formen auftreten, *Z. Math. Phys.* **27** (1882), 86–101.
4. A. JONQUIÈRE, Note sur la série $\sum_{n=1}^{\infty} x^n/n^s$, *Bull. Soc. Math. France* **17** (1889), 142–152.
5. V. A. LEBESGUE, Suite du Mémoire sur les applications du symbole $(a | b)$, *Jour. de Math.* **15** (1850), 215–237.
6. K. PRACHAR, "Primzahlverteilung." Springer-Verlag, Berlin/New York, 1957.
7. C. TRUESDELL, On a function which occurs in the theory of the structure of polymers, *Ann. Math.* **46** (1945), 144–157.